

# The Role of Sexual Abuse and Dysfunctional Attitudes in Perceived Stress and Negative Mood in Pregnant Adolescents: An Ecological Momentary Assessment Study



Kate Walsh PhD<sup>1,\*</sup>, Archana Basu PhD<sup>2</sup>, Catherine Monk PhD<sup>2,3,4</sup>

<sup>1</sup> Department of Epidemiology, Columbia University Medical Center, New York, New York

<sup>2</sup> Department of Psychiatry, Columbia University Medical Center, New York, New York

<sup>3</sup> Department of Psychiatry, Weill Medical College of Cornell University, New York, New York

<sup>4</sup> Department of Obstetrics and Gynecology, Columbia University Medical Center, New York, New York

## ABSTRACT

**Study Objective:** Latinas have the highest rates of adolescent pregnancy in the United States. Identifying means to improve the well-being of these young women is critical. The current study examined whether a history of child sexual abuse—itsself a risk factor for adolescent pregnancy—was associated with more perceived stress and negative mood over the course of pregnancy and whether dysfunctional attitudes explained these associations.

**Design and Setting:** This mixed methods study involved laboratory-based assessments of perceived stress, sexual abuse history, and dysfunctional attitudes, as well as Ecological Momentary Assessments (EMA) of mood states every 30 minutes during a 24-hour period once during each trimester of pregnancy.

**Participants:** Pregnant adolescents (N = 204, 85% Latina).

**Main Outcome Measures:** EMA mood states and laboratory-based retrospective self-reports of perceived stress.

**Results:** One in 4 pregnant adolescents had a history of sexual abuse. Sexually abused adolescents reported greater perceived stress during the first trimester relative to those without, though the groups did not differ on EMA negative mood ratings. Dysfunctional attitudes explained associations between sexual abuse and perceived stress. Sexual abuse was indirectly associated with the intercept and slope of negative mood through dysfunctional attitudes. Findings were circumscribed to sexual abuse and not other types of child abuse.

**Conclusions:** Identifying sexually abused pregnant adolescents and providing support and cognitive therapy to target dysfunctional beliefs may decrease stress during the first trimester as well as negative affect throughout pregnancy.

**Key Words:** Adolescent pregnancy, Child sexual abuse, Dysfunctional attitudes, Perceived stress, Negative mood, Ecological momentary assessment

## Introduction

Although adolescent pregnancy in the United States has declined significantly,<sup>1</sup> the United States continues to have the highest rate of all industrialized nations.<sup>2,3</sup> Rates of adolescent pregnancy are 3 times higher among Hispanic adolescents than among white adolescents,<sup>1</sup> and, independent of race and ethnicity, adolescent pregnancy is associated with increased risk for negative birth outcomes (eg, preterm birth and lower birth weight).<sup>4–6</sup> Identifying factors that may improve the health and well-being of Hispanic adolescent mothers-to-be has significant public health relevance.

Compared with their peers, pregnant adolescents are more likely to have a history of adverse childhood events,<sup>7</sup> including childhood sexual abuse (CSA).<sup>8</sup> Although CSA may increase risk for adolescent pregnancy, few studies have

examined how CSA impacts adolescents *during* pregnancy. According to traumagenic dynamics theory,<sup>9</sup> CSA could negatively impact pregnant adolescents by engendering negative thinking patterns or attitudes that increase risk for mood or anxiety disorders. CSA victims have negative attributions of events that are more internal, stable, and global than those of nonvictims,<sup>10</sup> and victims of more-severe CSA tend to endorse more self-blame attributions relative to victims of less-severe CSA.<sup>11,12</sup> CSA victims also report more dysfunctional beliefs and attitudes compared with those without CSA.<sup>13</sup> However, with the exception of a study documenting associations between CSA and depression during pregnancy,<sup>14</sup> the effects of potentially CSA-related dysfunctional beliefs and attitudes have not been studied among pregnant adolescents.

Dysfunctional beliefs and attitudes among adolescents with CSA histories could negatively impact pregnancy by increasing their stress and negative mood. Adult women with CSA histories evidence heightened negative mood and stress during pregnancy.<sup>15</sup> Pregnancy is an outcome of sexual relations and thus may activate fear networks associated with CSA exposure and increase the experience of stress, particularly for those with posttraumatic stress disorder (PTSD).<sup>15</sup> Several studies have linked CSA to the long-term development of disorders characterized by

The authors would like to acknowledge the following funding sources: Funded by National Institute on Drug Abuse (NIDA) with Grant # T32DA031099 (Dr Walsh), Sackler Institute Fellowship in Parent-Infant Studies from the Sackler Institute for Developmental Psychobiology (Dr Basu), and Funded by National Institute of Mental Health (NIMH) with Grant # MH077144 - 01A2 (Dr Monk). The authors indicate no conflicts of interest.

\* Address correspondence to: Kate Walsh, PhD, 722 W 168th St, New York, NY 10032

E-mail address: [klw2153@columbia.edu](mailto:klw2153@columbia.edu) (K. Walsh).

extreme dysphoria including borderline personality disorder, PTSD, and depression.<sup>16,17</sup> Of the few studies focused on pregnant adolescents, a cross-sectional study found that adolescents with CSA histories scored higher on general measures of stress.<sup>18</sup> A study of adult women found that while external stressors remained stable throughout pregnancy, pregnancy-related stress peaked during the first and third trimesters, and anxiety peaked during the third trimester.<sup>19</sup> Furthermore, among adult women assessed at 2 time points during pregnancy, perceived stress and anxiety were not associated with increased risk for preterm birth; however, those who exhibited an increase in stress and anxiety over the course of pregnancy had shorter gestation lengths.<sup>20</sup> Factors that contributed to increased stress over the course of the pregnancy were not explored in that study; findings suggest the importance of identifying malleable individual difference variables, including CSA history or dysfunctional attitudes, that could contribute to increases in stress or negative mood over the course of pregnancy.

### *The Current Study*

Although a burgeoning body of research has documented the impact of maternal stress during pregnancy on poor birth outcomes,<sup>21–24</sup> few studies have examined individual difference factors that could contribute to variability in maternal stress during pregnancy, and far fewer studies have focused on these associations in pregnant adolescents. A history of CSA is an individual difference factor that may elevate risk for psychological distress during pregnancy, perhaps through the development and maintenance of dysfunctional beliefs and attitudes that increase perceived stress and negative mood. The current study used data from a sample of pregnant, primarily Latina adolescents to examine whether a history of CSA was associated with more perceived stress and daily negative mood during repeated ecological momentary assessments (EMA) over the course of pregnancy. We collected data on a broad range of child maltreatment experiences as child maltreatment types are known to co-occur<sup>25</sup>; thus, we also tested whether the observed associations were circumscribed to CSA or whether they extended to other forms of maltreatment. We also examined whether dysfunctional beliefs and attitudes explained associations between CSA history and changes in stress and negative mood over the course of pregnancy. Compared with nonvictims, pregnant adolescents with CSA histories were expected to report more dysfunctional attitudes as well as greater perceived stress and increased negative mood initially and over the course of the prenatal period. It also was expected that dysfunctional attitudes would account for associations between CSA and trajectories of perceived stress and negative mood over time.

## **Materials and Methods**

### *Overall Design*

Participants were drawn from a prospective longitudinal observational study of pregnant adolescents measured at three time points: early (13–16 weeks), middle

(24–27 weeks), and late (34–37 weeks) pregnancy. Participants completed the perceived stress measure during a laboratory-based assessment. Over a 24-hour period, mood ratings were collected using EMA, which permits in-the-moment collection of data.

### *Participants*

Nulliparous pregnant adolescents, ages 14–19 and between 13–27 gestational weeks, were recruited through the Departments of Obstetrics and Gynecology at Columbia University Medical Center (CUMC) and Weill Cornell Medical College, and flyers posted in the CUMC vicinity. All adolescents had a healthy pregnancy at the time of recruitment. Participants were excluded if they smoked or used recreational drugs, lacked fluency in English, or were multiparous. Per a separate aim of the study, participants were excluded on the basis of chronic use of any of the following: nitrates, steroids, systemic migraine medications, stimulants, major and minor tranquilizers; and psychiatric medications. One pregnancy ended in fetal death; this participant was excluded from analysis. Participants were 204 pregnant adolescents (mean age = 17.9 years, SD = 1.2 years, range = 14–20 years). Approximately 85% (n = 173) were Latina; 48% (n = 98) were in the 12th grade, 25% (n = 51) were in the 11th grade, 11.8% (n = 24) were in the 10th grade, 8.3% (n = 17) were in the 9th grade, and 3.4% (n = 7) were in the 8th grade. Thirty-five percent (72 participants) reported a total annual family income of \$0–\$15,000, 34.3% (70 participants) reported \$16,000–\$25,000, 16.2% (33 participants) reported \$26,000–\$50,000, 2% (4 participants) reported \$51,000–\$100,000, and 0.5% (1 participant) reported greater than \$100,000. All participants provided written-informed consent, and all procedures were approved by the Institutional Review Board of the New York State Psychiatric Institute/CUMC.

### *Measures*

#### *Child Maltreatment History*

The Childhood Trauma Questionnaire (CTQ)<sup>26</sup> is a 28-item self-report questionnaire that retrospectively assesses 5 types of childhood maltreatment: emotional, physical, and sexual abuse and emotional and physical neglect. Each abuse type is assessed with 5 items measured on a Likert-type scale from 1 (never true) to 5 (very often true). Responses to the CTQ are summed, with higher scores indicating greater levels of each type of maltreatment. Numerous investigations attest to the reliability and validity of this measure.<sup>27,28</sup> The CTQ was administered once during pregnancy, typically during the first trimester, although 14.6% (30 participants) entered the study in their second trimester and received the CTQ during that session; victimization status according to the 5-item sexual abuse subscale was used for the primary analysis, and victimization status for any other maltreatment (physical or emotional abuse or physical or emotional neglect) was used for secondary analysis. Per cutoffs in the manual,<sup>26</sup> participants were considered CSA victims if they scored a 6 or higher on the sexual abuse subscale. Participants who

scored an 8 or higher on the physical abuse or neglect subscales, a 9 or higher on the emotional abuse subscale, and a 10 or higher on the emotional neglect subscale indicated victims of those forms of maltreatment.

#### *Dysfunctional Attitudes Scale (DAS)*<sup>29</sup>

The DAS is a 40-item measure of cognitive distortions that relate to or cause depression. DAS items are based on Beck's cognitive therapy model and present 7 major value systems: Approval, Love, Achievement, Perfectionism, Entitlement, Omnipotence, and Autonomy. A total score is obtained by summing the items. The DAS has good internal consistency, with  $\alpha$  values ranging from .84 to .92.<sup>29</sup> The DAS also has excellent stability, with test-retest correlations over 8 weeks of 0.80 to 0.84.<sup>29</sup> The DAS was administered once during pregnancy, typically during the first trimester, although 14.6% (30 participants) entered the study in their second trimester and received the DAS during that session.

#### *Perceived Stress Scale*<sup>30</sup>

The Perceived Stress Scale is a 14-item measure of the degree to which life situations during the previous month are perceived as stressful. Participants respond to items on a 5-point Likert scale ranging from 0 (never) to 4 (very often). The PSS was administered in the laboratory at the same three testing time points as EMA data were collected. Internal consistency reliability for the scale is 0.85, and the test-retest correlation is 0.85.<sup>30</sup>

#### *EMA Negative Mood*

A personal digital assistant was provided to participants to collect current mood states every 30 minutes for 24-hour periods timed to begin with each of the three lab assessments. Using a 5-point Likert scale (1 = low, 5 = high), participants rated 17 negative and positive mood states: cheerful, cooperative, responsible, caring, proud, friendly, productive, hard-working, lonely, nervous, angry, frustrated, competitive, strained, worried, irritated, and stressed. Participants provided ratings during waking hours each time an automated ambulatory blood pressure unit initiated a read (every 30 minutes). Participants were incentivized to provide ratings, earning \$0.10 per rating. As described elsewhere,<sup>31</sup> a composite negative mood score, which reflected experiencing events as straining or exceeding adaptive capacities and threatening well-being, was derived via factor analysis for each participant for each EMA session. The selected items were angry, frustrated, irritated, and stressed. A weight was created by dividing the number of diary entries at each session by the total number of diary entries across all study sessions. The average for the 4 items was calculated for each participant at each time point and multiplied by the weight at that session.

#### *Statistical Analyses*

Preliminary descriptive analyses and tests of bivariate associations were computed by using PASW Statistics Version 19.0 (IBM SPSS, Hong Kong, China). Tests of

hypotheses were conducted using Mplus Version 6.0 (Muthen & Muthen, Los Angeles, CA). Growth curve modeling was used to examine changes in the trajectories of perceived stress and negative mood during pregnancy. First, unconditional growth curve models were estimated separately by outcome to obtain intercept and slope fixed-effect estimates and to test for the presence of significant variance in these latent growth trajectory factors. Second, conditional growth curve models examined the influence of CSA and dysfunctional attitudes on the intercept and slope of perceived stress and negative mood during pregnancy. Data from only 3 time points (ie, once each trimester) were available for analysis; thus, nonlinear effects were not tested here. Time was parameterized such that the first trimester (T1) was at 0; subsequent time points were parameterized as months since T1. On average, 2.6 months (SD = 0.33 month) passed between T1 and T2, while 2.3 months (SD = 0.25 month) passed between T2 and T3. Model fit was considered acceptable if the  $\chi^2$  was nonsignificant, the comparative fit index (CFI) was 0.95 or greater, the root-mean-square error of approximation (RMSEA) was 0.06 or less, and the standardized root mean square residual (SRMR) was 0.08 or less.<sup>32,33</sup>

#### *Missing Data*

Adolescents who completed all 3 data collection sessions (61 participants) did not differ from those who completed 1 or 2 sessions (143 participants) on child abuse history,  $\chi^2(1,204) = 0.46, P = .54$ ; sexual abuse history,  $\chi^2(1,204) = 0.03, P = 1.0$ ; DAS,  $F(1,168) = 1.7, P = .20$ ; or PSS score at wave 1,  $F(1,125) = .004, P = .95$ , wave 2,  $F(1,124) = 3.00, P = .09$ , or wave 3,  $F(1,116) = 1.15, P = .29$ . However, adolescents who entered the study at wave 2 compared with wave 1 had greater perceived stress,  $F(1,124) = 6.49, P = .01$ . Furthermore, adolescents who missed 1 or 2 sessions had higher negative mood scores at wave 1,  $F(1,131) = 53.8, P < .001$ ; wave 2,  $F(1,124) = 29.9, P < .001$ ; and wave 3,  $F(1,118) = 19.9, P < .001$ , relative to adolescents who completed all 3 sessions. Missing data were handled via full information maximum likelihood (FIML) estimation procedures; however, because likelihood of missingness was associated with higher negative mood, sensitivity analyses were conducted just among those who completed all 3 sessions, to examine potential changes in findings.

## **Results**

### *Descriptive Statistics*

Correlations between study variables are presented in [Table 1](#). Approximately 25% (50 participants) reported a history of CSA. CSA victims did not differ from nonvictims in mean age,  $F(1,236) = 0.29, P = .59$ ; education,  $\chi^2(4,233) = 1.79, P = .78$ ; race/ethnicity,  $\chi^2(3,237) = 0.06, P = .81$ ; or family income,  $\chi^2(4,216) = 3.06, P = .55$ . Mean differences between CSA victims and nonvictims on dysfunctional attitudes, perceived stress, and negative mood are reported in [Table 2](#). Victims reported more dysfunctional attitudes and greater perceived stress during

**Table 1**  
Correlations Between Perceived Stress, Dysfunctional Attitudes, and Negative mood

Variable	1	2	3	4	5	6	7
1. T1 PSS	–	0.56 <sup>†</sup>	0.61 <sup>‡</sup>	0.40*	0.28 <sup>†</sup>	0.23*	0.01
2. T2 PSS		–	0.67 <sup>†</sup>	0.36	0.28*	0.29	0.32 <sup>†</sup>
3. T3 PSS			–	0.32	0.12	0.22	0.15
4. DAS total				–	0.33*	–0.02	0.33
5. Neg 1					–	0.12	–0.02
6. Neg 2						–	0.16
7. Neg 3							–

DAS, Dysfunctional Attitudes Scale; Neg, EMA negative mood; PSS, Cohen's Perceived Stress Scale total score.

\*  $P < .05$ .

†  $P < .01$ .

‡  $P < .001$ .

the first trimester relative to nonvictims. Victims did not report significantly more mean daily negative mood at each wave.

#### Growth Curve Models

Unstandardized intercept and slope estimates and variance from unconditional models are presented in Table 2. Models fit the data well as indicated by a nonsignificant  $\chi^2$ , CFI of 0.95 or greater, RMSEA of 0.05 or less, and SRMR of 0.08 or less. Significant fixed effects for the intercept and slope of perceived stress and negative mood were observed such that the intercept was significantly greater than 0 and the slope declined significantly over time for each variable.

#### Perceived Stress

Unstandardized intercept and slope estimates from conditional models predicting trajectories of perceived stress and negative mood also are presented in Table 3. For the model predicting perceived stress, model fit was good,  $\chi^2(df = 2) = 0.97$ ,  $P = .62$ , RMSEA  $< 0.001$ , CFI = 1.0, and SRMR = 0.03, and CSA predicted the intercept of perceived stress, although it did not predict the slope (see Fig. 1). When dysfunctional attitudes were added to the model, model fit also was good,  $\chi^2(df = 3) = 1.27$ ,  $P = .74$ , RMSEA  $< 0.001$ , CFI = 1.0, and SRMR = 0.03, and dysfunctional attitudes predicted perceived stress while CSA became nonsignificant (see Table 3). Neither was associated with the slope of perceived stress over the prenatal period. CSA was positively associated with dysfunctional attitudes (estimate = 12.06, SE = 4.69,  $P = .01$ ), and there was a significant indirect effect from CSA through dysfunctional attitudes to the intercept (estimate = 1.07, SE = .34,  $P = .013$ ) of perceived stress.

**Table 2**  
Raw Mean Differences Between CSA Victims and Nonvictims on Perceived Stress, Dysfunctional Attitudes, and Negative Mood

Measures (n)	CSA victim (n = 50)	Non-CSA victim (n = 134)	F (P value)
T1 PSS (118)	29.6 (6.6)	26.2 (6.4)	6.4 (.01)
T2 PSS (117)	28.4 (6.5)	25.8 (7.0)	3.4 (.07)
T3 PSS (107)	27.6 (4.9)	25.6 (5.6)	2.6 (.11)
DAS total (156)	122.8 (32.1)	106.2 (26.0)	11.4 (.001)
Neg 1 (122)	0.88 (.60)	0.83 (0.47)	0.25 (.62)
Neg 2 (117)	0.76 (.52)	0.70 (0.46)	0.33 (.56)
Neg 3 (107)	0.57 (.29)	0.53 (0.28)	0.43 (.51)

CSA, childhood sexual abuse; DAS, Dysfunctional Attitudes Scale; Neg, EMA negative mood; PSS, Cohen's Perceived Stress Scale total score.

#### Negative Mood

For the model predicting negative mood, model fit was good,  $\chi^2(df = 2) = 3.9$ ,  $P = .14$ , RMSEA = 0.07, CFI = 0.96, and SRMR = 0.05, but CSA was not associated with negative mood during pregnancy (see Fig. 1). When dysfunctional attitudes were added to the model, fit remained good,  $\chi^2(df = 3) = 4.6$ ,  $P = .20$ , RMSEA = 0.06, CFI = 0.95, and SRMR = 0.05, and dysfunctional attitudes were associated with increased negative mood at intercept, but with reduced negative mood over the prenatal period. Although CSA was not directly associated with negative mood trajectories, it was positively associated with dysfunctional attitudes (estimate = 12.49, SE = 4.67,  $P = .008$ ), and there was a significant indirect effect from CSA through dysfunctional attitudes to the intercept (estimate = 0.12, SE = 0.05,  $P = .02$ ) and slope (estimate = –0.02, SE = 0.01,  $P = .04$ ) of negative mood.

#### Secondary Analysis

To examine whether findings were specific to sexual abuse, secondary analyses were conducted with other forms of child abuse. Approximately 31% (64 participants) reported other child abuse; sexual abuse victims were excluded from this analysis. Victims of other child abuse did not differ significantly on age, ethnicity, education, or family income relative to nonvictims. Victims of other abuse did not have more dysfunctional attitudes, perceived stress, or negative mood at any wave; thus, growth models were not examined.

#### Sensitivity Analysis

Because individuals with higher negative mood at each time point were less likely to complete all 3 study sessions, analyses were conducted among those with complete data (61 participants) to examine whether parameter estimates changed. Consistent with findings from the larger sample, only dysfunctional attitudes contributed to the intercept of perceived stress (estimate = 0.08, SE = 0.04,  $P = .02$ ), and CSA indirectly affected the intercept of perceived stress through dysfunctional attitudes (estimate = 1.73, SE = 0.84,  $P = .04$ ). However, the associations between dysfunctional attitudes and negative mood that were evident in the larger sample were not significant among complete cases.

## Discussion

The current study is one of the first to examine how CSA impacts perceived stress and negative mood during pregnancy in a sample of pregnant, primarily Latina adolescents who comprise the largest ethnic/racial group of pregnant adolescents in the United States. This study also examined whether dysfunctional beliefs account for associations between CSA and perceived stress and negative mood during pregnancy. One in 4 pregnant adolescents reported a history of CSA, which is lower than the 45% of pregnant adolescents who report CSA according to a recent meta-analysis.<sup>8</sup> Growth curve models revealed that perceived stress and negative mood both declined over the course of pregnancy; however, changes were minimal. A history of CSA was not associated with EMA-assessed negative mood, but it was

**Table 3**  
Conditional and Unconditional Model Results

Measure	PSS int (Variance)	PSS Slope (Variance)	Neg int (Variance)	Neg Slope (Variance)
Unconditional model	27.6 <sup>†</sup> (29.4 <sup>†</sup> )	−0.26* (0.30)	0.96 <sup>†</sup> (0.33 <sup>†</sup> )	−0.08 <sup>†</sup> (0.01 <sup>†</sup> )
Conditional models	PSS int (SE)	PSS slope (SE)	Neg int (SE)	Neg slope (SE)
Model 1: CSA	2.38 (1.1)*	−0.23 (0.20)	0.03 (0.09)	0.06 (0.10)
Model 2: CSA	1.44 (1.0)	−0.24 (0.21)	−0.03 (0.11)	0.02 (0.02)
Model 2: DAS total	0.09 (0.02) <sup>†</sup>	−0.001 (0.004)	0.007 (0.002) <sup>†</sup>	−0.001 (0.001)*


CSA, childhood sexual abuse; DAS, Dysfunctional Attitudes Scale; Neg, EMA negative mood; Int, Intercept; PSS, Cohen's Perceived Stress Scale total score. Nonstandardized parameters are presented: \* $P < .01$ , <sup>†</sup> $P < .001$ , <sup>‡</sup> $P < .05$ .

related to increased perceived stress during the first trimester and dysfunctional attitudes mediated this association. Although CSA did not directly relate to negative mood during pregnancy, a history of CSA was indirectly associated with higher initial negative mood and a decrease in negative mood over the course of pregnancy that operated through dysfunctional attitudes. Consistent with hypotheses and traumagenic dynamics theory, which was developed to explicate the impact of sexual abuse,<sup>9</sup> secondary analyses indicated that findings are specific to sexual abuse and not exposure to child abuse more generally.

Consistent with prior work,<sup>15</sup> we found heightened perceived stress during pregnancy among women with CSA histories. This effect was specific to the first trimester, which also is consistent with other studies.<sup>18</sup> Studies with adults typically have found perceived stress to increase during the first and third trimesters (eg, Da Costa et al<sup>19</sup>); however, pregnant adolescents in the current study demonstrated slight linear decreases in perceived stress, which is congruent with at least 1 study finding a linear decline in psychological distress over the course of pregnancy.<sup>34</sup>

We also found a novel association between dysfunctional attitudes and higher levels of perceived stress and negative mood during the first trimester. Although changes in perceived stress and negative mood were quite small over pregnancy, adolescents evidenced a decline in both outcomes and greater dysfunctional attitudes were associated with a swifter decline in negative mood over pregnancy. Different findings for perceived stress and negative mood may stem from methodological differences in how

perceived stress was measured in the laboratory as a retrospective recall of previous month events, while negative mood was assessed in the moment several times during a 24-hour period. As further support for construct differences, low-level positive correlations emerged for trimesters 1 and 2, but the constructs were not significantly correlated by trimester 3. Although results on dysfunctional attitudes and negative mood could reflect regression to the mean, they also could reflect habituation to the negative cognitions and emotions that pregnancy may have initially triggered given that pregnancy is a result of sexual activity. Furthermore, because adolescents with more dysfunctional beliefs demonstrated swifter decreases in negative mood over the course of the study relative to those with fewer dysfunctional beliefs, adolescents with CSA histories who initially had more dysfunctional beliefs may have experienced psychological integration, and possibly aspects of resolution, through an emerging change in identity (ie, motherhood).<sup>35</sup> Specifically, attachment schemas may become salient during pregnancy,<sup>36</sup> and the opportunity to form a new attachment bond as a caregiver may have a positive effect on levels of perceived stress and negative mood over the course of pregnancy. Families in which CSA occurs are often characterized as dysfunctional.<sup>37</sup> Participants for the current study were involved in a study involving regular contact with health care professionals who were monitoring their health and well-being; as a result, they may have had access to better services and care than they otherwise might have received. For adolescents with CSA histories, it is possible that this increased support



**Fig. 1.** Changes in perceived stress and negative mood during pregnancy by childhood sexual abuse (CSA) history. Neg mood, EMA negative mood; PSS, Cohen's Perceived Stress scale; t1-t3, trimesters 1-3.

made a substantial difference in their mood states during this potentially high-stress period.

Results should be interpreted in the context of limitations. First, the sample was small and, due to other study aims, only high-functioning pregnant adolescents (eg, those without substance use) were eligible for participation. There is ample evidence that pregnant adolescents with sexual abuse histories are more likely to use substances during pregnancy (eg, Stevens-Simon and McAnarney,<sup>18</sup> Bayatpour et al<sup>38</sup>); thus, excluding these individuals may have restricted our ability to examine the most severely abused adolescents. Second, adolescents were not specifically asked whether the index pregnancy was the result of CSA; such experiences could be associated with greater negative affect or stress during pregnancy. Third, individuals with higher negative mood ratings were less likely to complete all of the study assessments, and sensitivity analyses among those with complete data indicated no associations between dysfunctional attitudes and negative mood ratings. Had girls with higher negative mood completed all assessments, associations may have differed.

The current study highlights a number of directions for further research. While this study focused on experiences during pregnancy, future studies should examine how CSA, dysfunctional attitudes and beliefs, perceived stress, and negative mood are associated with birth outcomes and early child development. Examining maternal mental health diagnoses (eg, depression, PTSD) during pregnancy may illuminate whether particular symptoms have differential relationships with perceived stress and negative mood. Efforts should be made to improve study retention among girls with increased levels of negative mood.

This study has important clinical implications. Rates of adolescent pregnancy are higher among Latinas; thus, it is critical to understand methods to improve their health and experiences (ie, level of stress, mood) during pregnancy. Adolescents with CSA histories had increased dysfunctional attitudes and perceived stress during the first trimester when exposure to toxins and stressors could have a particularly detrimental effect on fetal development.<sup>39,40</sup> Identifying adolescents with CSA histories and dysfunctional attitudes early in pregnancy and providing increased support as well as cognitive therapy to reduce dysfunctional attitudes could result in potentially beneficial reduction in perceived stress and negative mood in this group.

## References

- Ventura SJ, Curtin SC, Abma JC, et al: Estimated pregnancy rates and rates of pregnancy outcomes for the United States, 1990–2008. *Natl Vital Stat Rep* 2012; 60:1
- Darroch JE, Singh S, Frost JJ: Differences in teenage pregnancy rates among five developed countries: the roles of sexual activity and contraceptive use. *Fam Plann Perspect* 2001; 33:244
- Hamilton BE, Ventura SJ: Birth Rates for US teenagers Reach Historic Lows for All Age and Ethnic Groups. Atlanta, GA, US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics, 2012
- Chen XK, Wen SW, Fleming N, et al: Teenage pregnancy and adverse birth outcomes: a large population based retrospective cohort study. *Int J Epidemiol* 2007; 36:368
- Gortzak-Uzan L, Hallak M, Press F, et al: Teenage pregnancy: risk factors for adverse perinatal outcome. *J Matern Fetal Med* 2001; 10:393
- Klein JD: American Academy of Pediatrics Committee on Adolescent Pregnancy: Adolescent pregnancy: current trends and issues. *Pediatrics* 2005; 116:281
- Hillis SD, Anda RF, Dube SR, et al: The association between adverse childhood experiences and adolescent pregnancy, long-term psychosocial consequences, and fetal death. *Pediatrics* 2004; 113:320
- Noll JG, Shenk CE, Putnam KT: Childhood sexual abuse and adolescent pregnancy: A meta-analytic update. *J Pediatr Psychol* 2009; 34:366
- Finkelhor D, Browne A: The traumatic impact of child sexual abuse: a conceptualization. *Am J Orthop* 1985; 55:530
- Weninger K, Ehlers A: Dysfunctional cognitions and adult psychological functioning in child sexual abuse survivors. *J Trauma Stress* 1998; 11:281
- Owens GP, Chard KM: Cognitive distortions among women reporting childhood sexual abuse. *J Interpers Violence* 2001; 16:178
- Quas JA, Goodman GS, Jones DP: Predictors of attributions of self-blame and internalizing behavior problems in sexually abused children. *J Child Psychol Psychiatry* 2003; 44:723
- Henderson D, Hargreaves I, Gregory S, et al: Autobiographical memory and emotion in a non-clinical sample of women with and without a reported history of childhood sexual abuse. *Br J Clin Psychol* 2002; 41:129
- Benedict MI, Paine LL, Paine LA, et al: The association of childhood sexual abuse with depressive symptoms during pregnancy, and selected pregnancy outcomes. *Child Abuse Negl* 1999; 23:659
- Lev-Wiesel R, Daphna-Tekoa S: Prenatal posttraumatic stress symptomatology in pregnant survivors of childhood sexual abuse: a brief report. *J Loss Trauma* 2007; 12:145
- Green JG, McLaughlin KA, Berglund PA, et al: Childhood adversities and adult psychiatric disorders in the National Comorbidity Survey Replication I: associations with first onset of DSM-IV disorders. *Arch Gen Psychiatry* 2010; 67:113
- Saunders BE, Kilpatrick DG, Hanson RF, et al: Prevalence, case characteristics, and long-term psychological correlates of child rape among women: a national survey. *Child Maltreatment* 1999; 4:187
- Stevens-Simon C, McAnarney ER: Childhood victimization: relationship to adolescent pregnancy outcome. *Child Abuse Negl* 1994; 18:569
- Da Costa D, Larouche J, Dritsa M, et al: Variations in stress levels over the course of pregnancy: factors associated with elevated hassles, state anxiety and pregnancy-specific stress. *J Psychosom Res* 1999; 47:609
- Glynn LM, Schetter CD, Hobel CJ, et al: Pattern of perceived stress and anxiety in pregnancy predicts preterm birth. *Health Psychol* 2008; 27:43
- Dole N, Savitz DA, Hertz-Picciotto I, et al: Maternal stress and preterm birth. *Am J Epidemiol* 2003; 157:14
- Goldenberg RL, Culhane JF, Iams JD, et al: Epidemiology and causes of preterm birth. *Lancet* 2008; 371:75
- Rondo P, Ferreira R, Nogueira F, et al: Maternal psychological stress and distress as predictors of low birth weight, prematurity and intrauterine growth retardation. *Eur J Clin Nutr* 2003; 57:266
- Wadhwa PD, Entringer S, Buss C, et al: The contribution of maternal stress to preterm birth: issues and considerations. *Clin Perinatol* 2011; 38:351
- Clemmons JC, Walsh K, DiLillo D, et al: Unique and combined contributions of multiple child abuse types and abuse severity to adult trauma symptomatology. *Child Maltreat* 2007; 12:172
- Bernstein D, Fink L: Manual for the Childhood Trauma Questionnaire. New York, NY, The Psychological Corporation, 1998
- Bernstein DP, Stein JA, Newcomb MD, et al: Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl* 2003; 27:169
- Scher CD, Stein MB, Asmundson GJ, et al: The childhood trauma questionnaire in a community sample: psychometric properties and normative data. *J Trauma Stress* 2001; 14:843
- Weissman AN: The Dysfunctional Attitude Scale: a Validation Study. Dissertation Abstracts International. ProQuest Information Learning, Pennsylvania, US, University of Pennsylvania, 1979, pp 1389–1390
- Cohen S, Kamarck T, Mermelstein R: A global measure of perceived stress. *J Health Soc Behav* 1983;385
- Spicer J, Werner E, Zhao Y, et al: Ambulatory assessments of psychological and peripheral stress-markers predict birth outcomes in teen pregnancy. *J Psychosom Res* 2013; 75:305
- Browne MW, Cudeck R, Bollen KA, et al: Alternative ways of assessing model fit. *Sage Focus Editions* 1993; 154:136
- Lt Hu, Bentler PM: Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equat Model* 1999; 6:1
- Entringer S, Buss C, Shirtcliff EA, et al: Attenuation of maternal psychophysiological stress responses and the maternal cortisol awakening response over the course of human pregnancy. *Stress* 2010; 13:258
- Ammaniti M, Baumgartner E, Candelori C, et al: Representations and narratives during pregnancy. *Infant Ment Health J* 1992; 13:167
- Monk C, Leight KL, Fang Y: The relationship between women's attachment style and perinatal mood disturbance: implications for screening and treatment. *Arch Womens Ment Health* 2008; 11:117
- Bhandari S, Winter D, Messer D, et al: Family characteristics and long-term effects of childhood sexual abuse. *Br J Clin Psychol* 2011; 50:435
- Bayatpour M, Wells RD, Holford S: Physical and sexual abuse as predictors of substance use and suicide among pregnant teenagers. *J Adolesc Health* 1992; 13:128
- Sandman CA, Davis EP, Buss C, et al: Prenatal programming of human neurological function. *Int J Pept* 2011; 2011:837596
- Sandman CA, Davis EP, Buss C, et al: Exposure to prenatal psychological stress exerts programming influences on the mother and her fetus. *Neuroendocrinology* 2012; 95:8